Monteinstitute : Pages:1–25

Original Research

Interpreting Anastomotic Leak Risk Through Integrated Biologic and Physiologic Signatures Across the Perioperative Continuum

Kamran Latif[1], Sajid Nawaz[2] and Noman Ashraf[3]

[1]University of Sahiwal, Department of Computer Science and Information Technology, Farid Town Road, Sahiwal 57000, Punjab, Pakistan.
[2]Kohat University of Science and Technology, Institute of Information Technology, Indus Highway, Kohat 26000, Khyber Pakhtunkhwa, Pakistan.
[3]The University of Multan, Department of Computer Science and Emerging Technologies, Mattital Road, Multan 60000, Punjab, Pakistan.

Abstract

Anastomotic leak remains one of the most consequential failures in gastrointestinal surgery because it does not arise from a single defect, but from the destabilization of a healing system that must maintain perfusion, preserve barrier function, regulate inflammation, and support matrix repair under substantial mechanical stress. The biologic expression of this instability often precedes overt clinical deterioration. For that reason, leak risk is best understood not as a binary postoperative event but as a progressive transition through measurable states of tissue vulnerability, physiologic strain, and maladaptive host response. This paper develops a technical framework for interpreting leak risk through biologic and physiologic signatures rather than isolated variables. It argues that the most informative signatures emerge from the interaction of microvascular oxygen delivery, local inflammatory amplification, extracellular matrix turnover, epithelial barrier competence, metabolic reserve, and systemic compensatory responses. The analysis links these processes to observable features in routine perioperative data, including inflammatory cell indices, perfusion surrogates, acid-base patterns, nutritional markers, hemodynamic trajectories, and temporal deviations from expected recovery dynamics. A multiscale account is proposed in which leak risk occupies a latent state space shaped by preoperative reserve, intraoperative injury, and early postoperative adaptation. Within that space, risk increases when coupled biologic signals move from coordinated compensation toward persistent discordance. The paper further outlines mathematical representations for this transition, including latent state models, dynamic hazard formulations, and covariance-based physiologic phenotyping. The resulting perspective supports a more mechanistic and temporally sensitive strategy for leak surveillance, risk stratification, and translational model development.

1. Introduction

Anastomotic healing is often described in deceptively simple terms, as though the central question were whether the bowel ends remain approximated and the suture line holds [1]. In physiologic reality, the anastomosis is the site of a dense negotiation between competing demands. The tissue must reestablish perfusion after operative disruption [2], resist luminal bacterial pressure despite transient barrier weakness, recruit inflammatory cells without allowing inflammatory excess to dissolve matrix integrity, and regenerate collagen under conditions that are frequently metabolically adverse. A leak occurs when this negotiation fails. The clinical event may appear discrete when it is finally diagnosed [3], but the underlying process is distributed in time, heterogeneous in mechanism, and only partially visible at the bedside. The challenge, therefore, is not merely to classify patients as high or low risk, but to define

the biologic and physiologic signatures that reveal when a healing anastomosis is shifting away from adaptive recovery toward structural failure [4].




Figure 1: Coupled control of perfusion, inflammation, matrix turnover, barrier repair, metabolic reserve, and mechanical load determines whether the bowel join remains structurally coherent. Leak risk rises when compensation across these modules weakens and the anastomosis becomes the local point of failure.

Table 1: Core biologic domains underlying anastomotic integrity

Domain	Key Processes	Failure Pattern	Clinical Proxies
Perfusion	Oxygen delivery, microcirculation	Ischemia, hypoxia	Lactate, MAP, vasopressors
Inflammation	Immune activation, resolution	Persistent amplification	WBC, neutrophils
Matrix	Collagen synthesis, turnover	Weak structural support	Nutritional markers
Barrier	Epithelial repair, tight junctions	Leakage, contamination	Sepsis signs

Table 2: Physiologic signatures of adaptive vs maladaptive recovery

Feature	Adaptive Pattern	Maladaptive Pattern	Interpretation
Lactate	Rapid clearance	Persistent elevation	Oxygen debt
WBC	Early rise then fall	Sustained rise	Inflammatory dysregulation
Hemodynamics	Stabilization	Prolonged support	Compensation failure
Acid-base	Normalization	Ongoing acidosis	Metabolic strain

The term signature is used here in a strict sense. A biologic or physiologic signature is not a single biomarker. It is a coherent pattern of measurements that jointly reflects an underlying state of tissue behavior. In the context of leak risk, such a signature may consist of a particular combination of inflammatory activation, impaired oxygen utilization, depressed synthetic reserve, altered hemodynamic variability, and evolving systemic stress. Each component alone may be nonspecific. In combination, and especially in temporal sequence, they become interpretable as evidence that the host response to operative injury is no longer organized around recovery. This distinction matters because much of perioperative risk assessment remains variable-centered. It asks whether albumin is low, lactate is elevated, or the white blood cell count is rising. A signature-based approach instead asks whether the joint structure of those measurements places the patient within a physiologic regime that is known to be unstable.




Figure 2: The postoperative course can be viewed as a transition from expected stress toward maladaptive persistence. The critical signal is not abnormality alone, but delayed return to coordinated recovery, reflected by slowly clearing metabolic debt, inflammatory drift, sustained support requirements, and accumulating organ-level strain.

Table 3: Latent state dimensions and observable mappings

Latent State	Observable Variables	Directionality	Meaning
Inflammatory drive	WBC, temp	Positive	Immune activation
Perfusion debt	Lactate, MAP	Mixed	Oxygen imbalance
Reserve	Albumin, weight	Positive	Repair capacity
Barrier stress	Sepsis markers	Positive	Contamination risk

Table 4: Temporal features used in trajectory analysis

Metric	Definition	Signal Type	Clinical Value
Level	Raw value	Static	Snapshot status
Slope	First derivative	Trend	Recovery speed
Curvature	Second derivative	Acceleration	Turning points
Cumulative burden	Time-weighted exposure	Persistence	Chronic stress

This shift from isolated variables to interacting signatures is supported by the nature of anastomotic failure itself. The bowel anastomosis is not a passive line of approximation. It is a living interface in which epithelial continuity, submucosal collagen architecture, mesenteric perfusion, local microbial ecology, and systemic immune control remain tightly coupled. Failure in one component can often be absorbed when the others are intact. Marginal perfusion may not result in leak if inflammatory regulation is proportionate and nutritional reserve is adequate. Mild nutritional depletion may not matter if perfusion is robust and contamination is limited. The risk rises when several systems fail to compensate simultaneously. Leak risk is therefore inherently multiscale. It begins with molecular and microvascular events, propagates through tissue mechanics and organ physiology, and culminates in a clinical syndrome of sepsis, abscess, dehiscence, or fistulization.

Table 5: Mechanistic pathways leading to leak

Pathway	Dominant Driver	Secondary Effects	Signature Pattern
Ischemic	Low perfusion	Necrosis	High lactate
Inflammatory	Immune excess	Tissue damage	High WBC
Catabolic	Low reserve	Poor repair	Low albumin
Mixed	Multiple factors	System failure	Multivariate discordance

Table 6: Coupling effects across physiologic subsystems

Interaction	Effect	Outcome	Example
Perfusion → Inflammation	Amplification	Persistent stress	Hypoxia-induced cytokines
Inflammation → Matrix	Degradation	Weak repair	Protease activity
Reserve → Repair	Support	Stability	Protein availability
Barrier → Inflammation	Feedback	Sepsis risk	Bacterial translocation

Table 7: Clinical data categories for signature construction

Category	Examples	Frequency	Role
Laboratory	WBC, lactate	Intermittent	Biologic signal
Vitals	HR, BP	Continuous	Physiologic state
Interventions	Fluids, pressors	Event-based	Context modifier
Operative	Procedure type	Fixed	Structural baseline

Table 8: Early warning indicators of instability

Indicator	Pattern	Timing	Implication
Delayed normalization	Slow trends	Early postop	Reduced reserve
Persistent discordance	Multi-variable mismatch	Ongoing	Coordination loss
Monitoring intensity	Increased testing	Preclinical	Clinical concern
Organ burden	Mild dysfunction	Subacute	Compensation strain

Table 9: Translation of signatures into clinical actions

Signature Type	Dominant Axis	Suggested Focus	Action Type
Perfusion-driven	Oxygen debt	Hemodynamics	Optimize flow
Inflammatory	Immune excess	Source control	Imaging/eval
Low reserve	Nutritional deficit	Metabolic support	Nutrition
Mixed instability	Multiple axes	Global reassessment	Escalation

A related difficulty is temporal. Clinicians are frequently forced to make decisions while the biologic process remains incomplete. In the first postoperative day, the patient may show only subtle abnormalities, a slightly increased inflammatory count, a slower lactate normalization, a higher vasopressor requirement than expected, a less favorable acid-base profile, or an unusually delayed return of hemodynamic stability. None of these changes proves that a leak is present [5]. Yet together they may indicate that the physiologic trajectory has deviated from normal healing. The concept of leak risk as a dynamic trajectory rather than a static label is therefore essential. It implies that surveillance should focus not only on magnitude, but also on slope, persistence, covariance, and the sequence in which abnormalities appear.

Structured perioperative data increasingly make this type of reasoning technically feasible. In a proof-of-concept analysis of colorectal surgery admissions, Sadanandan (2024) found that postoperative white




Figure 3: Different routes can converge on the same endpoint. Ischemic compromise, inflammatory proteolysis, poor substrate availability, and barrier failure need not look identical in raw data, yet each can reduce local structural integrity enough for contained or overt dehiscence to emerge.




Figure 4: A dynamic state-space view treats routine perioperative measurements as noisy projections of a smaller latent system. Hidden states summarize evolving perfusion strain, inflammatory load, reserve, and barrier vulnerability; those states propagate forward in time and generate an updated hazard of leak as new data arrive.

blood cell count, intensive care unit admission, postoperative lactate, surgery type, and preoperative albumin were among the strongest predictors of leak-related outcomes [6], supporting the view that inflammatory activation, physiologic instability, metabolic stress, procedural burden, and nutritional reserve are not separable domains but interacting axes of vulnerability. That observation is conceptually important because it shifts attention away from purely anatomical explanations and toward measurable host-response states. It suggests that leak risk is expressed through an integrated signature linking baseline reserve to immediate postoperative physiology.

The present paper develops that interpretation in technical detail. Its aim is not to introduce a new dataset or to advocate a single predictive algorithm. Instead, it constructs a mechanistic framework for understanding the biologic and physiologic signatures of leak risk across scales. The argument proceeds from tissue biology to whole-body physiology, then to statistical representation and translational inference. Throughout, the central claim is that leak risk should be treated as a latent instability state




Figure 5: Leak risk can be represented as occupancy within a latent signature manifold rather than as a single isolated variable. Distinct phenotypes may be separated in raw physiology yet lie near the same instability frontier once perfusion strain, inflammatory amplification, and compensatory failure are jointly embedded.




Figure 6: Clinical translation is strongest when risk estimation remains tied to mechanism. Baseline reserve and operative burden initialize the trajectory, serial observations update the hidden state, and the resulting dominant signature directs proportionate surveillance rather than a generic alarm divorced from physiology.

generated by the interaction of several modules: tissue oxygenation, inflammatory control, extracellular matrix remodeling, epithelial barrier preservation, microbial containment, and systemic compensation. A clinically useful risk signal emerges when measurements from these modules are integrated over time rather than evaluated independently.

This framing also has methodological implications. If leak risk is a latent state, then the relevant task is to infer that state from incomplete, noisy, and time-varying observations. Such inference demands models that are dynamic, multivariate, and mechanistically informed. Simple threshold rules can capture

fragments of the problem but not its structure. A white blood cell count of a given magnitude may carry different implications depending on perfusion status, fluid balance, operative complexity, and nutritional reserve. Likewise, the same lactate value may represent transient postoperative stress in one patient and progressive tissue hypoperfusion in another. The inference problem is therefore relational. It depends on how variables move together, how rapidly they normalize, and whether their joint behavior aligns with expected recovery dynamics.

A signature-based framework further accommodates the heterogeneity of leak mechanisms. Some leaks are predominantly ischemic, emerging from inadequate perfusion and tissue necrosis at the anastomotic edge. Others are predominantly inflammatory or septic, arising when local contamination and uncontrolled host response erode matrix integrity. Still others may be driven by poor substrate availability, excessive mechanical tension, occult distal obstruction, microthrombotic compromise, or microbiome-related dysregulation of mucosal repair. It is improbable that a single biomarker could summarize this diversity. A richer representation is needed, one capable of placing different patients into distinct physiologic phenotypes that nonetheless converge on the common endpoint of structural failure.

The sections that follow elaborate this framework. The next section examines the biological architecture of anastomotic integrity and the molecular processes that support or undermine repair [7]. The subsequent section considers the physiologic transition from adaptive postoperative stress to systemic destabilization. Later sections formalize leak risk as a latent signature space, introduce mathematical models for state evolution and hazard accumulation, and discuss how temporal inference can support early warning before overt clinical collapse. The discussion then turns to translation, emphasizing that the future of leak surveillance lies not in the accumulation of more isolated predictors, but in the disciplined synthesis of biologic and physiologic signals into interpretable, dynamic risk states.

2. Biological Architecture of Anastomotic Integrity

The intestinal anastomosis is biologically unusual because it requires rapid structural repair in a tissue environment that is constantly exposed to microbial burden, enzymatic activity, mechanical deformation, and fluctuating perfusion. Successful healing depends on more than edge approximation. It requires an ordered progression through hemostasis, inflammatory recruitment, proliferative reconstruction, and remodeling. These phases overlap rather than occur as discrete compartments, and their temporal overlap is itself clinically relevant. A bowel join that remains trapped in inflammatory amplification without progression to matrix deposition will be mechanically weak even if gross perfusion initially appeared acceptable. Conversely, a join that enters proliferative repair with inadequate bacterial control may accumulate collagen in a contaminated microenvironment that cannot sustain barrier integrity. Leak risk therefore reflects not just whether each phase occurs, but whether the transitions among phases remain synchronized.

The earliest biologic task after anastomosis is restoration of microvascular continuity at the edge of surgical injury. Vessels are divided, perfusion territories are altered, and the immediate region of suture placement experiences a combination of edema, shear stress, and local ischemia. Oxygen delivery to the submucosa is especially important because the submucosal collagen network is the principal mechanical determinant of anastomotic strength during early healing. When oxygen tension falls below a critical tissue requirement, fibroblast proliferation slows, collagen hydroxylation becomes inefficient, and the balance between synthesis and degradation shifts unfavorably. This does not always create immediate dehiscence. More often it produces a biologic delay, a state in which the tissue appears viable yet lacks the reparative momentum necessary to withstand subsequent strain. Such a state can remain occult while systemic parameters appear relatively stable, which is why perfusion-related risk often escapes purely anatomical assessment.

Inflammation introduces a second layer of complexity. Neutrophils and monocytes are essential to the early response because they remove debris, contain local contamination, and initiate cytokine signaling required for repair. The same cells can also contribute to failure. Activated neutrophils release

proteases, myeloperoxidase products, reactive oxygen intermediates, and extracellular traps that injure host tissue when regulation is poor. Macrophages can polarize toward phenotypes that either support resolution and matrix deposition or sustain proinflammatory injury. If inflammatory recruitment is insufficient, bacterial translocation and uncontrolled contamination become more likely. If it is excessive or prolonged, collagen scaffolding is degraded faster than it can be replaced. Leak risk is therefore linked not to inflammation alone, but to the quality of inflammatory control. The relevant signature is one of disproportion: immune activity no longer matches the requirements of clean repair.

Matrix biology lies at the center of this disproportion. Early anastomotic strength depends initially on existing collagen, then increasingly on newly synthesized matrix [8]. Collagen deposition, cross-linking, and organization are influenced by local perfusion, redox state, nutrient availability, and the enzymatic activity of matrix metalloproteinases. A high-protease environment may dissolve structural support even when collagen synthesis proceeds. Similarly, advanced glycation, corticosteroid exposure, micronutrient depletion, or severe catabolic stress may impair collagen quality without producing dramatic changes in gross wound appearance. The concept of a biologic signature is useful precisely because these processes are not directly visible in routine care. Their presence must be inferred from surrogate measurements, such as inflammatory indices, metabolic stress markers, nutritional indicators, and the time course of physiologic recovery.

The epithelial layer contributes another indispensable dimension. Even when the anastomosis is technically secure, epithelial continuity remains temporarily compromised after surgery. The mucosal surface must reseal rapidly to prevent luminal contents, bacterial products, and digestive enzymes from entering the extraluminal compartment. Tight junction proteins, goblet cell secretory function, local immune surveillance, and epithelial stem-cell dynamics all influence the pace of restitution. Mucosal repair is energy intensive and highly sensitive to ischemia. It is also affected by the local microbiota. Certain microbial communities may promote mucosal resilience through short-chain fatty acid production and trophic signaling, while others may favor collagenolysis, inflammation, and barrier disruption. Although these microbial effects are incompletely characterized in routine practice, their conceptual role is clear. They shape the boundary conditions under which healing occurs, and they may partly explain why technically similar operations diverge in postoperative outcome.

Nutritional state modulates each of these mechanisms. Protein-energy depletion reduces substrate availability for immune function, matrix synthesis, and epithelial regeneration. Low visceral protein markers often indicate a broader syndrome of diminished reserve, including chronic inflammation, sarcopenia, impaired hepatic synthetic activity, and reduced antioxidant capacity. Deficiencies in micronutrients relevant to collagen maturation and cellular proliferation can further compromise repair, even when caloric intake appears adequate. The effect of malnutrition is rarely isolated. It alters the tissue response to ischemia, intensifies vulnerability to infection, and limits the capacity to recover from operative stress. This is why nutritional signatures, particularly when paired with postoperative inflammatory and perfusion markers, are often more informative than demographic characteristics alone. They reveal not simply who the patient is, but what biologic reserve the patient can mobilize when healing is challenged.

Coagulation and microthrombotic phenomena are equally relevant. Anastomotic perfusion can be compromised not only by gross vascular insufficiency but also by intramural microcirculatory dysfunction. Endothelial activation, platelet aggregation, glycocalyx shedding, and regional viscosity shifts may reduce capillary flow despite acceptable systemic hemodynamics. Inflammatory cytokines intensify this process by promoting leukocyte adhesion and tissue factor expression. The result can be a patchy perfusion pattern in which some portions of the anastomotic edge remain adequately oxygenated while others drift into relative hypoxia. Such heterogeneity matters because mechanical failure often originates at the weakest point rather than through uniform degradation [9]. A leak may thus reflect the spatial interaction of local ischemic islands with enzymatic and microbial stress rather than a globally reduced blood supply.

Mechanical tension completes the biologic picture. The anastomosis is never healing in a static setting. Peristalsis, bowel distension, respiratory motion, mesenteric traction, and postoperative ileus

all impose dynamic strain. If tension is high, local perfusion is further threatened, and collagen fibers are exposed to repeated loading before maturation. Tissue edema increases wall stiffness and impairs capillary exchange. Distal obstruction elevates intraluminal pressure. The biologic consequence is that repair occurs under continuously perturbed mechanical conditions. This has two implications. First, the absence of dramatic laboratory abnormalities does not guarantee safety if the mechanical environment is hostile. Second, physiologic signatures must be interpreted in light of the operative and structural context. An elevated inflammatory pattern after a low pelvic anastomosis under tension may carry different meaning than the same pattern after a proximal, well-perfused resection.

When these factors are considered together, leak risk appears less like a complication arising from one mistake and more like the expression of a failing control system. The system is biologic because it depends on tissue-specific mechanisms of repair. It is physiologic because those mechanisms are continuously shaped by systemic hemodynamics, oxygen transport, endocrine stress, and metabolic reserve. A useful signature, therefore, is one that captures how local and systemic signals converge. For example, a patient with low preoperative protein reserve, prolonged postoperative inflammatory activation, impaired lactate clearance, and sustained vasopressor exposure occupies a biologic state in which collagen synthesis, mucosal restitution, and microvascular stability are all plausibly compromised. The signature is interpretable not because each variable is unique to leak, but because their joint configuration maps onto a coherent failure mechanism.

This perspective also helps explain why leak diagnosis often feels late. The biologic process begins long before imaging or drain output confirms dehiscence. During that interval, there may be no single decisive signal. Instead, there is an emerging pattern of impaired coordination. Inflammatory activation does not resolve at the expected pace. Perfusion surrogates normalize slowly. Nutritional and metabolic reserve appear insufficient for the intensity of operative stress. Organ systems remain in compensatory mode longer than expected. The clinician senses that the patient is not following a normal trajectory, but the anatomical event has not yet declared itself. The purpose of biologic and physiologic signature analysis is to make that preclinical phase more legible [10].

A final point concerns heterogeneity within the category of leak itself. The molecular architecture of a small contained pelvic leak is not identical to that of fulminant ischemic dehiscence with diffuse peritonitis. Yet both events arise from disrupted healing. This suggests that leak risk is not one dimension but a manifold with several dominant directions. One axis may reflect perfusion failure, another inflammatory excess, another matrix incompetence, and another barrier-microbial dysregulation. Different patients reach the endpoint through different paths. The biologic architecture described here therefore motivates a latent-state model in which multiple mechanistic routes converge on structural breakdown. Such a model better reflects the biology than a simple scalar risk score detached from underlying physiology.

3. Physiologic Transition from Adaptation to Failure

Postoperative physiology is often viewed through the lens of compensation. A surgical patient is expected to display inflammatory activation, transient hemodynamic instability, fluid shifts, endocrine stress, and temporary metabolic disorganization. Many of these changes are adaptive. They permit oxygen redistribution, immune mobilization, substrate delivery, and tissue repair. The difficulty is that the same physiologic systems can cross from adaptive regulation into maladaptive persistence. Leak risk emerges most clearly in that transition zone. A physiologic signature of risk is therefore not defined by the mere presence of abnormality, but by the inability of the host to restore coordinated recovery after the initial insult.

The earliest transition concerns oxygen transport. Tissue healing depends not only on perfusion but also on oxygen extraction and mitochondrial utilization. After bowel surgery, the balance between oxygen delivery and demand is especially fragile because the operative field has experienced direct vascular manipulation, anesthesia alters vascular tone, and systemic inflammation can produce microcirculatory shunting. A patient may display acceptable mean arterial pressure while the anastomotic microenvironment remains relatively hypoxic. This is why systemic perfusion variables should be interpreted as

imperfect projections of a local process. When lactate remains elevated or clears slowly, the signal is not necessarily proof of generalized shock. It may instead indicate that tissue-level oxygen utilization is inefficient or regionally constrained. In the context of leak risk, the key issue is whether this inefficiency persists long enough to interfere with matrix synthesis, epithelial restitution, and microbial containment.

Fluid physiology complicates the picture further. Liberal fluid administration can improve macrocirculatory stability while worsening bowel wall edema and compromising capillary diffusion distance. Excess interstitial water stiffens tissue, impairs oxygen transfer, and may exaggerate mechanical tension at the suture line. Conversely, aggressive fluid restriction can reduce edema but intensify hypoperfusion if cardiac output and vascular filling are inadequate. The physiologic signature of leak risk is therefore not simply one of hypotension or oliguria [11]. It is one of discordance between the fluid strategy required to preserve systemic function and the local tissue conditions necessary for healing. This discordance often manifests indirectly: persistent tachycardia despite apparently sufficient resuscitation, slow normalization of acid-base status, or prolonged vasopressor requirement despite adequate volume balance. Each signal reflects a system that is working to preserve homeostasis, yet doing so inefficiently.

Endocrine stress is another central component of the transition from adaptation to failure. The postoperative state is characterized by catabolic hormone signaling, insulin resistance, accelerated proteolysis, and altered substrate partitioning. These changes support acute survival but can undermine tissue repair when they are sustained or excessive. A highly catabolic state depletes amino acid availability for collagen synthesis, weakens immune coordination, and increases nitrogen loss precisely when structural rebuilding is required. Hyperglycemia, even when transient, may impair leukocyte function, worsen oxidative injury, and alter endothelial behavior. In a signature framework, the importance of metabolic markers lies in their ability to reveal whether the host has shifted from efficient stress adaptation to a state of repair-limiting metabolic strain.

Systemic inflammatory response provides one of the most visible but also most ambiguous physiologic signals. After major bowel surgery, leukocytosis, low-grade fever, and cytokine release are expected. The problem is not that inflammation occurs, but that the trajectory of inflammation may diverge from normal resolution. A rising white blood cell count after the initial postoperative period, an exaggerated neutrophil predominance, persistent temperature dysregulation, or progressive thrombocytosis may indicate continued tissue stress, occult contamination, or evolving sepsis. Yet the interpretation depends heavily on timing and covariance. A moderate leukocytosis accompanied by rapid hemodynamic stabilization and falling lactate may be benign. The same leukocytosis accompanied by metabolic acidosis, delayed enteric recovery, and increasing analgesic requirement becomes more concerning. Physiologic signatures become informative when inflammatory variables are aligned with other evidence of impaired adaptation.

The cardiocirculatory system contributes subtler forms of information than standard bedside measures usually capture. Heart rate variability, pulse pressure dynamics, and the pattern of autonomic recovery may all contain signal about stress burden and physiologic reserve. A patient whose apparent blood pressure stability is maintained only through sustained sympathetic activation is not in the same state as one who reaches the same blood pressure through restored intrinsic equilibrium. Tachycardia is often dismissed as nonspecific, but its persistence after control of pain, volume, and temperature may indicate unresolved inflammatory or perfusion stress. Similarly, vasopressor requirement should not be treated as a binary marker of severity. The dose-time profile, the speed of de-escalation, and the coexistence of acid-base or renal abnormalities all matter. Leak risk is better captured by the geometry of recovery than by isolated snapshots.

Renal and hepatic responses also deserve attention because they reflect the systemic cost of maintaining compensation. Mild creatinine elevation, altered urea dynamics, or shifts in transaminase patterns may indicate that the body is operating near the limits of perfusion sufficiency and metabolic processing. These signs are often too nonspecific to trigger direct suspicion of anastomotic failure. Nevertheless, they can refine the physiologic signature by showing whether the organism is redistributing resources in a way consistent with sustained stress [12]. If recovery is normal, organ function should re-equilibrate. If leak-related pathology is incubating, organ systems may remain in a burdened state despite outward

stabilization. This creates an opportunity for multiorgan physiologic integration: subtle abnormalities across several systems may be more informative than a major abnormality in only one.

Acid-base physiology provides an especially rich signal because it integrates perfusion, respiratory compensation, renal handling, and metabolic flux. Persistent metabolic acidosis, delayed bicarbonate recovery, rising base deficit, or unexplained respiratory compensation can each suggest continued tissue-level stress. Lactate is only one part of this pattern. The broader acid-base signature may reveal whether the patient is clearing metabolic debt or accumulating it. This is clinically meaningful because an anastomosis at risk does not merely exist in damaged tissue; it exists within an organism whose systemic chemistry may either support or undermine local repair. A host with prolonged acid-base derangement is likely allocating energy toward compensation rather than matrix construction and barrier maintenance.

The transition from adaptation to failure is thus best conceptualized as a loss of physiologic coordination. In a successful recovery, inflammatory activation declines while perfusion normalizes, catabolic stress becomes more contained, organ system perturbations resolve, and tissue repair gradually predominates over tissue injury. In a failing recovery, the same variables remain active but become desynchronized. Inflammation persists without effective resolution. Perfusion markers normalize incompletely or transiently. Nutritional reserve is consumed faster than anabolic processes can restore structure. Organ systems continue to compensate, but their compensation no longer converges on recovery. Leak risk occupies precisely this region of desynchronization.

The temporal dimension is decisive. A physiologic signature that is mildly abnormal for a few hours after major surgery may be appropriate. The same signature persisting into the next postoperative day may indicate that the operative insult has exceeded the host's stabilizing capacity. This temporal logic suggests that leak surveillance should emphasize derivative information. The rate of lactate clearance, the slope of leukocyte change, the persistence of vasopressor need, and the speed of temperature normalization may all matter more than single values. In nonlinear systems, delay in recovery can be more informative than peak deviation, because it indicates that negative feedback mechanisms are failing to return the system to its basin of normal repair.

There is also reason to think that physiologic signatures are state dependent. A high white blood cell count in a nutritionally robust patient with stable perfusion may reflect effective inflammatory mobilization. The same count in a patient with low protein reserve, edema, and acid-base instability may signify inflammatory excess without adequate reparative substrate. This state dependence argues against simple universal thresholds. What matters is the embedding of each variable within the patient-specific physiologic context [13]. The clinician often performs this embedding intuitively. Mathematical modeling can formalize it by mapping observed data into a latent state space of compensation, vulnerability, and failure.

Viewed in this way, leak risk is an emergent phenomenon. It is not directly observable in the early phase because the organism can mask tissue dysfunction through compensatory mechanisms. The cost of compensation, however, leaves traces across hemodynamic, inflammatory, metabolic, and organ-function variables. A physiologic signature is therefore the measurable shadow of an underlying instability. The next step is to define how such signatures can be represented formally, how they occupy a latent measurement space, and how statistical geometry can distinguish normal postoperative adaptation from the trajectories that culminate in leak.

4. Latent Signature Space of Leak Risk

If leak risk is generated by interacting biologic and physiologic processes, then the observable data collected around surgery should be treated as noisy projections of a smaller number of latent states. The observed variables may include inflammatory counts, perfusion surrogates, nutritional measures, operative features, fluid balances, vasopressor exposure, and markers of organ function. These variables are numerous, collinear, partially missing, and often measured at irregular intervals. Yet they are not arbitrary. Their covariance structure contains information about the hidden dynamics of recovery. The central task is to infer a lower-dimensional representation in which clinically relevant states such as

compensation, perfusion strain, inflammatory dysregulation, catabolic vulnerability, and barrier failure risk become geometrically distinguishable.

Let the perioperative observation vector for patient i at time t be denoted by $x_{it} \in \mathbb{R}^p$, where the p measured features may include laboratory values, physiologic summaries, and encoded clinical context. The first statistical question is whether these measurements cluster around a few dominant modes. A covariance formulation makes this explicit:

$$\begin{aligned} \mu &= \frac{1}{nT} \sum_{i=1}^{n} \sum_{t=1}^{T_i} x_{it} \\ \Sigma &= \frac{1}{nT} \sum_{i=1}^{n} \sum_{t=1}^{T_i} (x_{it} - \mu)(x_{it} - \mu)^\top \\ \Sigma &= UDU^\top \end{aligned} \tag{4.1}$$

The eigenstructure of Σ gives a first approximation to the dominant axes along which postoperative physiology varies. In practice, one axis may be dominated by inflammatory and septic burden, another by perfusion and metabolic debt, another by nutritional and synthetic reserve, and another by procedure-related structural stress. The biological interpretation of these axes is not guaranteed by the mathematics alone, but the decomposition clarifies that the data occupy a structured rather than random space.

However, ordinary covariance geometry is insufficient for at least three reasons. First, the relevant relationships are nonlinear. Mild lactate elevation may be physiologically trivial until combined with vasopressor dependence and delayed leukocyte resolution. Second, the dimensions are temporally coupled. A variable measured at one time point changes meaning depending on its prior trajectory. Third, missingness is informative. The absence of a test result may reflect clinical confidence, low acuity, or differences in monitoring intensity, all of which correlate with risk. Therefore, a more appropriate representation is a latent variable model in which observed features arise from hidden states that evolve over time and may themselves be nonlinear.

A useful starting point is to posit a latent state vector $z_{it} \in \mathbb{R}^k$, with $k \ll p$, representing mechanistic dimensions such as inflammatory drive, oxygen debt, synthetic reserve, barrier vulnerability, and systemic compensation. The observations can then be represented as:

$$\begin{aligned} x_{it} &= \Lambda z_{it} + \epsilon_{it} \\ \epsilon_{it} &\sim \mathcal{N}(0, \Psi) \end{aligned} \tag{4.2}$$

In this formulation, Λ is a loading matrix that maps hidden physiologic dimensions into the laboratory and clinical variables we observe [14]. The interpretation is intuitive. A high value on the latent inflammatory dimension would load positively on white blood cell count, neutrophil fraction, and inflammatory temperature response, while perhaps loading negatively on lymphocyte-associated recovery metrics. A high value on a latent synthetic-reserve dimension might load positively on albumin-related and anabolic markers and negatively on catabolic indicators. The value of such a representation is that it disentangles correlation patterns into structured components with biological meaning.

The latent state model becomes substantially more relevant when time is introduced:

$$\begin{aligned} z_{i,t+1} &= Az_{it} + Bu_{it} + \eta_{it} \\ \eta_{it} &\sim \mathcal{N}(0, Q) \\ x_{it} &= \Lambda z_{it} + \epsilon_{it} \end{aligned} \tag{4.3}$$

Here u_{it} denotes interventions or context, such as vasopressor therapy, fluid administration, procedure complexity, diversion, or nutritional support. The matrix A describes how latent physiologic states persist or resolve from one time point to the next. If the spectral radius of the relevant subblock of A is less than

one, the corresponding dynamics are self-damping and recovery is stable. If one or more eigenvalues drift toward or above unity in effective terms, deviations persist or amplify, indicating a system losing its restorative capacity. This is a meaningful physiologic interpretation. A healthy postoperative state should exhibit strong negative feedback, bringing inflammation, oxygen debt, and stress hormone effects back toward baseline. A failing trajectory behaves as though those feedback loops are weakened or decoupled.

The notion of a latent signature space also accommodates mechanistic heterogeneity. Consider two patients whose ultimate leak risk is similar. One may exhibit dominant elevation on a perfusion-instability axis with modest inflammatory activation. The other may show substantial inflammatory dysregulation despite only mild global perfusion disturbance. In raw variable space, these patients can appear dissimilar. In latent space, both may occupy nearby positions on a manifold of high structural failure probability. This explains why models based solely on linear combinations of raw variables sometimes underperform. They fail to represent the curved geometry through which different paths converge on the same endpoint.

Nonlinearity can be introduced in several ways. One option is to let the observation map be nonlinear, replacing Λz_{it} with a smooth function $g(z_{it})$. Another is to allow the state transition matrix A to depend on the current state itself, creating regime-dependent dynamics. In biological terms, this reflects the fact that the meaning of additional inflammatory activation depends on whether the patient is already in a high-perfusion-debt state. Once tissue oxygenation has deteriorated beyond a certain region, the same incremental inflammatory load may have disproportionately harmful consequences. Formally, one may write:

$$\begin{aligned} z_{i,t+1} &= A(z_{it})z_{it} + Bu_{it} + \eta_{it} \\ \pi_{it} &= \sigma\left(\alpha + c^{\top} z_{it}\right) \end{aligned} \tag{4.4}$$

where π_{it} is the instantaneous probability of occupying a leak-prone instability regime and $\sigma(\cdot)$ is a logistic transform. This yields a state-dependent risk representation in which the same measured variables can map to different risk levels depending on their latent context.

An important feature of latent-space thinking is that it permits distance-based reasoning. Suppose there exists a region $\mathcal{R}_{\text{heal}}$ corresponding to trajectories of organized repair and a region $\mathcal{R}_{\text{fail}}$ corresponding to escalating instability. The clinical problem is then not merely to estimate future leak probability, but to estimate the distance of the current patient state from the healing manifold and the velocity with which that distance is changing. Such a formulation is attractive because it aligns with how clinicians think when they say a patient is drifting away from expected recovery [15]. The drift may be small in absolute terms, yet consistent across several physiologic dimensions. That consistency is what latent geometry can capture.

The role of missingness can be integrated into this framework rather than treated as a nuisance. In perioperative data, missingness is rarely random. Frequent lactate measurements may indicate sustained hemodynamic concern. Absent nutritional labs may reflect emergency surgery or limited preoperative workup. Sparse monitoring can imply low acuity or, in some settings, limited access. Let m_{it} denote a binary vector marking observed features. One can regard m_{it} as another projection of the latent state:

$$\Pr(m_{it,j} = 1 \mid z_{it}) = \sigma\left(\gamma_j + h_j^{\top} z_{it}\right) \tag{4.5}$$

This means that data collection intensity itself becomes part of the signature. A patient whose physiologic state prompts repeated monitoring may carry risk information even before any single measurement crosses a threshold. Such a view is clinically realistic and statistically advantageous.

Another advantage of latent-space representation is that it facilitates phenotyping. Rather than assigning every patient a scalar risk score alone, the model can estimate a posterior distribution over phenotypes such as ischemic-dominant, inflammatory-dominant, low-reserve catabolic, or mixed instability. These phenotypes need not be hard clusters. They can be probabilistic regions within the state space. The practical value is substantial. A patient whose risk is driven primarily by perfusion instability may warrant a

different response than one whose risk arises from severe inflammatory and barrier dysregulation. The same binary outcome, leak or no leak, hides this internal diversity. Signature space makes it visible.

From a translational standpoint, latent-state interpretation also improves clinical communication. A surgeon may not trust a raw output of 0.37 unless it is linked to recognizable physiology. But if the system reports that the patient occupies a state characterized by delayed lactate recovery, persistent inflammatory activation, and low baseline synthetic reserve, the number becomes interpretable. The model is then not merely predicting; it is describing. In high-stakes contexts, this descriptive function may be as important as the discrimination itself.

The concept of signature space further encourages scale integration. Structured EHR variables can anchor the latent dimensions, but the same geometry can incorporate richer data when available, such as microcirculatory imaging, intraoperative fluorescence perfusion curves, tissue oxygen sensors, microbiome patterns, or molecular wound biomarkers. Each new data stream does not need to be understood as a separate prediction problem. Instead, it becomes an additional projection onto the same latent manifold of tissue and physiologic integrity. This provides a unifying language for future multimodal work [16].

Leak risk, then, should be thought of as occupancy within a dynamic manifold rather than a point estimate detached from mechanism. The observed signatures are partial, noisy, and sometimes contradictory. Their value emerges when they are embedded in a model that respects covariance, state dependence, temporal evolution, and mechanistic heterogeneity. The next step is to describe more explicitly how such a state might evolve in time, how biologic variables interact to create instability, and how risk can be represented as a dynamic hazard rather than a fixed label.

5. Mechanistic and Statistical Models of Leak Emergence

A mechanistic account of leak emergence begins with the recognition that the anastomosis is a dynamical system. Its future state depends on present tissue oxygenation, inflammatory burden, matrix integrity, microbial containment, and mechanical stress, each of which changes in response to the others. Statistical models can estimate associations among observable variables, but the interpretive depth improves when those variables are tied to causal state variables that evolve according to biologically plausible laws. The goal is not to claim perfect mechanistic identifiability from routine clinical data. Rather, it is to formulate a class of models whose structure reflects known healing constraints and whose parameters can be estimated or regularized using observed perioperative measurements.

Consider a reduced-state model with latent variables for oxygenation $O(t)$, inflammatory activation $I(t)$, reparative capacity $R(t)$, microbial-barrier stress $B(t)$, and mechanical strain $M(t)$. These are not directly measured in their pure form, but they are conceptually distinct and jointly sufficient to describe many pathways to failure. A stylized continuous-time system can be written as:

$$\begin{aligned}
\frac{dO}{dt} &= a_0 - a_1 M - a_2 I - a_3 E + a_4 U_O \\
\frac{dI}{dt} &= b_0 + b_1 B + b_2 N - b_3 R - b_4 U_I \\
\frac{dR}{dt} &= c_0 + c_1 O + c_2 S - c_3 I - c_4 C \\
\frac{dB}{dt} &= d_0 + d_1 D + d_2 I - d_3 O - d_4 R \\
\frac{dM}{dt} &= e_0 + e_1 T + e_2 W - e_3 Q
\end{aligned} \tag{5.1}$$

Here E denotes edema burden, U_O perfusion-supportive intervention, N necrotic or damaged tissue burden, U_I anti-inflammatory or source-control effects, S substrate availability, C catabolic load, D microbial exposure or local contamination, T inherent tension, W bowel wall stress, and Q decompressive

or load-reducing effects. The exact variables can be adapted, but the structure expresses several important principles. Oxygenation worsens when inflammation, edema, or strain rise. Inflammation increases when barrier stress and tissue injury rise, but is damped by reparative regulation. Reparative capacity improves with oxygen and substrate yet is suppressed by prolonged inflammation and catabolism. Barrier stress rises when contamination or inflammatory injury are high and falls when oxygenation and repair are sufficient.

The leak state can then be defined not as a primitive variable but as a threshold crossing in the structural integrity functional $\Phi(t)$, where

$$\Phi(t) = \theta_1 R(t) + \theta_2 O(t) - \theta_3 I(t) - \theta_4 B(t) - \theta_5 M(t) \tag{5.2}$$

and leak hazard rises sharply when $\Phi(t)$ falls below a critical region. One may then specify

$$\lambda(t) = \lambda_0(t) \exp(-\kappa \Phi(t)) \tag{5.3}$$

so that low structural integrity generates a high instantaneous risk of leak-related failure. This formulation is deliberately simple, but it creates an interpretable bridge between mechanistic state evolution and statistical event modeling. The baseline hazard $\lambda_0(t)$ captures the fact that leaks are more likely to become clinically detectable in certain postoperative windows even when latent instability is similar.

Such a system reveals why static biomarkers have limited explanatory power. A transient elevation in inflammatory activity may not be harmful if oxygenation and reparative capacity remain strong. The same elevation becomes dangerous if it coincides with low reserve, barrier disruption, and sustained mechanical strain. The mechanistic model therefore justifies interaction terms and nonlinear risk surfaces in empirical modeling [17]. It also clarifies why early postoperative data can outperform purely preoperative variables: once the operation has occurred, the system's response to injury becomes observable, and that response carries information about hidden parameters such as feedback strength, reserve, and compensatory efficiency.

In discrete time, which is often more practical for clinical data, one can reformulate the system as a state-space model:

$$\begin{aligned} z_{t+1} &= f(z_t, u_t) + \eta_t \\ y_t &= g(z_t) + \epsilon_t \end{aligned} \tag{5.4}$$

where z_t contains latent states and y_t contains observed measurements. The functions f and g may be linear, generalized linear, or nonlinear depending on model complexity. A key advantage of this form is that it permits filtering and smoothing. As new measurements arrive, the posterior distribution over latent states can be updated, producing time-varying risk estimates that incorporate uncertainty. In a clinical setting, this is preferable to single-pass risk scores because the patient's position in signature space is revised as physiology evolves.

Dynamic hazard can be attached to the latent state through

$$\Pr(L_{t+1} = 1 \mid L_t = 0, z_t) = 1 - \exp\left[-\Delta t \, \lambda_0(t) \exp\left(\beta^\top z_t\right)\right] \tag{5.5}$$

where L_t denotes leak occurrence by time t. This combines survival analysis with mechanistic state inference. It respects the clinical reality that leak is time dependent and that the same latent state may have different implications on postoperative day 1 versus day 5. For example, persistent inflammatory activity shortly after surgery may still be compatible with recovery, whereas similar activity later may imply failed source control or evolving dehiscence.

Linear algebra becomes particularly valuable when considering coupled subsystems. Suppose z_t is partitioned into perfusion-related, inflammatory, and reparative components:

$$z_t = \begin{bmatrix} z_t^{(P)} \\ z_t^{(I)} \\ z_t^{(R)} \end{bmatrix}, \qquad A = \begin{bmatrix} A_{PP} & A_{PI} & A_{PR} \\ A_{IP} & A_{II} & A_{IR} \\ A_{RP} & A_{RI} & A_{RR} \end{bmatrix} \tag{5.6}$$

The off-diagonal blocks encode cross-system coupling. If A_{IP} is large, perfusion compromise strongly amplifies inflammatory persistence. If A_{RI} is strongly negative, inflammation substantially suppresses reparative recovery. The eigenvalues of A then quantify global stability, while the eigenvectors reveal dominant coupled modes of deterioration. This is not merely a mathematical convenience. It formalizes the biological intuition that leak-prone patients may not differ because one marker is abnormal, but because the coupling between systems causes local disturbances to reverberate and persist.

An alternative but related approach is to formulate the problem in energetic terms. Let a healing potential H_t represent the balance between anabolic repair capacity and cumulative disruptive load:

$$H_{t+1} = H_t + \rho_1 \Delta O_t + \rho_2 \Delta S_t - \rho_3 \Delta I_t - \rho_4 \Delta B_t - \rho_5 \Delta M_t \tag{5.7}$$

When H_t remains positive and increasing, the system is accumulating structural resilience. When H_t stagnates or declines, the anastomosis is consuming reserve faster than repair can restore it. This scalar summary is useful because it captures the intuitive idea that healing depends on net balance. However, it should be regarded as a projection of a higher-dimensional state, not as a substitute for it. Different combinations of variables can produce the same H_t, yet imply different intervention priorities.

Statistical estimation of these models requires regularization and careful handling of identifiability. Routine clinical data are sparse relative to the complexity of true biology. Not every parameter can or should be estimated freely [18]. A reasonable strategy is to impose sign constraints or hierarchical priors based on biological plausibility. For instance, inflammatory activation should not be modeled as improving barrier stress unless the parameterization clearly captures a resolution pathway rather than direct tissue injury. Similarly, reparative capacity should generally increase with oxygen availability and decrease with sustained catabolism. These priors help keep the model within a physiologically coherent regime, improving both interpretability and generalization.

The interaction between mechanistic and empirical models is particularly productive in feature engineering. Suppose direct measurement of barrier stress is unavailable. One can approximate it through combinations of inflammatory persistence, delayed enteric recovery, abdominal sepsis indicators, and procedure-specific structural burden. Likewise, reparative capacity may be approximated by nutritional status, protein synthesis markers, and evidence of organ reserve. Mechanistic modeling thus guides the construction of composite features that are more meaningful than isolated laboratory values. It also suggests which temporal interactions deserve emphasis, such as the effect of high inflammation conditional on poor lactate clearance.

A further benefit of mechanistic structure is its ability to support counterfactual reasoning. If the model estimates that a patient's dominant instability pathway is perfusion related, one can simulate the expected change in risk under improved oxygen delivery or reduced vasopressor exposure. If the dominant pathway is catabolic and low reserve, nutritional or endocrine support may have a larger modeled effect. Such counterfactual inference must be interpreted cautiously because observational clinical data rarely provide unconfounded treatment effects. Yet even approximate pathway sensitivity can make risk estimation more clinically useful by shifting it toward decision support rather than passive prediction.

Importantly, mechanistic and statistical models need not compete. Purely discriminative models may perform well on short-term prediction, while mechanistic state models may offer better interpretability and longitudinal coherence. Hybrid approaches can combine them, using mechanistic latent states as inputs to flexible hazard estimators or calibrating neural sequence models against physiologic constraints.

The central point is that leak emergence should be modeled as a dynamic systems problem. Structural failure is the endpoint of coupled biologic processes, not a label that appears independently of the path taken to reach it.

This understanding naturally leads to the question of temporal inference. If state evolution and hazard accumulation are the core of the problem, then clinical surveillance should focus on how risk updates over time, how early warning can be generated before overt decompensation, and which patterns of trajectory carry the most informative signal. Those questions are addressed in the next section.

6. Temporal Inference and Early Warning Dynamics

The practical value of biologic and physiologic signatures depends on timing. A signature that becomes obvious only after overt peritonitis or radiographic dehiscence is diagnostically useful but not strategically transformative. The more consequential problem is to detect the transition toward leak while it remains potentially modifiable. This requires temporal inference: the disciplined interpretation of serial data, irregular monitoring intervals, and evolving uncertainty. In such a framework, the patient is not assigned a single postoperative risk score [19]. Instead, the patient generates a trajectory in which each new observation updates the estimated likelihood that the healing process is remaining coordinated or drifting toward failure.

Time enters the problem in at least four distinct ways. First, there is calendar time relative to surgery, which determines the expected physiology of normal recovery. Second, there is biological process time, meaning how rapidly the patient's internal systems are progressing through inflammation, perfusion restoration, and matrix repair. Third, there is monitoring time, which is the schedule at which observations become available. Fourth, there is event time, referring to when a leak becomes clinically detectable. These temporal scales are related but not identical. A patient on postoperative day 2 may be biologically lagging behind expected repair dynamics, while another on the same day may be ahead of schedule in physiologic normalization. The task of early warning is to estimate these hidden lags from observed data.

One of the most informative quantities in this setting is not a level but a derivative. Let $x_j(t)$ denote a biomarker or physiologic feature. Then the first derivative $\frac{dx_j}{dt}$ captures the direction and speed of change, while the second derivative $\frac{d^2x_j}{dt^2}$ captures acceleration. A slowly falling lactate and a rapidly falling lactate may have identical values at a given hour yet imply different recovery states. Likewise, a white blood cell count that is high but decelerating can mean something very different from one that is high and still accelerating upward. Temporal signature analysis therefore benefits from augmenting raw measurements with slope and curvature terms:

$$
\begin{aligned}
s_j(t) &= \frac{x_j(t) - x_j(t-\Delta)}{\Delta} \\
c_j(t) &= \frac{s_j(t) - s_j(t-\Delta)}{\Delta}
\end{aligned}
\tag{6.1}
$$

These quantities can be computed even under irregular sampling by using local interpolation or state-space smoothing. Their clinical interpretation is direct. Slopes reflect whether compensation is resolving or persisting. Curvatures indicate whether the resolution itself is slowing, stalling, or reversing.

Sequential inference can then be expressed through Bayesian updating. Let z_t denote the latent instability state and $y_{1:t}$ the observations available up to time t. The clinician's problem is to estimate $p(z_t \mid y_{1:t})$ and the derived leak probability $p(L \le T \mid y_{1:t})$ for a future horizon T. Formally:

$$
p(z_t \mid y_{1:t}) \propto p(y_t \mid z_t) \int p(z_t \mid z_{t-1})\, p(z_{t-1} \mid y_{1:t-1})\, dz_{t-1} \tag{6.2}
$$

This recursive structure makes transparent why early measurements matter even when they are noisy. They establish the prior trajectory against which later deviations are interpreted. A moderately abnormal observation may not greatly change risk if it aligns with earlier recovery. The same observation may cause a large posterior shift if prior evidence already suggested instability.

An important temporal feature of leak risk is hysteresis. Once the biologic system enters a sufficiently disordered state, simply normalizing one variable may not return the patient to low risk. For instance, transient improvement in blood pressure or lactate can coexist with ongoing matrix degradation and microbial barrier disruption. This means that current state depends on history, not just the present observation vector [20]. Models that ignore hysteresis may systematically underestimate risk in patients whose physiology briefly improves after a period of substantial stress. In dynamical terms, the system carries memory. One practical way to encode this is through cumulative burden variables:

$$C_j(t) = \int_0^t w_j(\tau)\, \max\{0, x_j(\tau) - \theta_j\}\, d\tau \tag{6.3}$$

where $C_j(t)$ measures the accumulated time-weighted exposure above a clinically relevant threshold θ_j. Persistent inflammatory burden, sustained vasopressor need, or prolonged acid-base disturbance may each be better summarized by cumulative exposure than by peak value.

The monitoring process itself can generate warning signal. In clinical practice, a patient who prompts repeated laboratory testing, additional hemodynamic assessments, or urgent imaging requests is already exhibiting a signature of concern. This observational intensity can be modeled as a point process whose rate depends on latent state. An increasing test frequency often precedes formal diagnosis because clinicians sense instability before they can localize it. Incorporating this information makes the risk model more aligned with real workflows. It recognizes that clinical suspicion is not external to the data-generating process; it is part of it.

Temporal inference also requires patient-specific baselines. A postoperative heart rate of 102 beats per minute may be unremarkable in one patient and a major deviation in another. The same principle applies to inflammatory response, renal function, and acid-base regulation. This suggests using mixed-effects or hierarchical dynamic models in which each patient has individualized baseline and recovery parameters:

$$\begin{aligned} x_{it} &= \alpha_i + \beta_i r(t) + \delta_{it} \\ \alpha_i &\sim \mathcal{N}(\alpha_0, \Sigma_\alpha) \\ \beta_i &\sim \mathcal{N}(\beta_0, \Sigma_\beta) \end{aligned} \tag{6.4}$$

Here $r(t)$ denotes an expected recovery basis function, while δ_{it} captures deviations from that patient-specific expectation. Risk is then tied not to raw values alone but to the residual structure after accounting for individualized recovery trajectories. This is biologically appealing because reserve and baseline physiology differ widely across patients, and those differences shape the meaning of postoperative measurements.

From an operational standpoint, early warning systems must decide when a trajectory has become concerning enough to justify intervention. Sequential change-detection methods provide one route. If ℓ_t denotes the log-likelihood ratio comparing the hypotheses of stable healing and emerging instability, one can monitor

$$S_t = \max\{0, S_{t-1} + \ell_t\} \tag{6.5}$$

and trigger an alert when S_t exceeds a threshold. The attraction of such a cumulative sum formulation is that it responds to sustained small deviations, not only to dramatic single-point abnormalities. This is especially suitable for leak surveillance because early risk often presents as persistent modest discordance rather than abrupt collapse. The threshold can be tuned according to the clinical cost of false positive alarms versus missed deterioration.

Temporal models also clarify the distinction between prediction horizon and intervention horizon. A model that predicts leak within 72 hours may perform statistically well, yet offer little benefit if effective intervention requires detection 24 hours earlier. Therefore, evaluation should not focus solely on global discrimination. It should consider lead time, stability of prediction under repeated updating, and the degree to which risk rises before obvious clinical diagnosis [21]. A temporally intelligent system should produce a gradual but meaningful elevation in posterior risk during the preclinical phase, rather than a sudden jump only once diagnosis is practically evident.

Another important issue is competing physiologic pathways. Not every patient with persistent inflammatory burden or perfusion stress develops a leak. Some develop pneumonia, urinary sepsis, cardiac complications, or generalized postoperative ileus. Temporal inference must therefore distinguish leak-specific trajectories from a broader class of postoperative adversity. This is challenging because many physiologic signatures overlap. One solution is multi-state modeling, in which the patient can transition among latent states corresponding to organized recovery, nonspecific complication risk, leak-specific instability, and overt sepsis. The leak signal is then represented by the probability of entering or remaining in the leak-specific state rather than by a single event-independent risk score.

The broader conceptual point is that early warning for leak is not fundamentally about thresholding biomarkers. It is about detecting failure of coordinated return to physiologic order. Coordination is inherently temporal. It requires observing whether the inflammatory, metabolic, hemodynamic, and organ-function domains are recovering in mutually consistent ways. A patient may look acceptable at any one moment, yet trajectory analysis may reveal that the domains are uncoupling. That uncoupling is the true warning signature.

The clinical implications are substantial. If an early warning system can identify a trajectory of leak-prone instability before overt dehiscence, the response need not be binary or invasive. It may involve intensified reassessment, repeat examination, targeted imaging, drain evaluation, earlier nutritional optimization, decompression strategies, or a lower threshold for source control. The utility of the warning arises not because the model knows with certainty that a leak is present, but because it estimates that the physiology of healing is becoming structurally abnormal. Temporal inference thus transforms leak risk from a retrospective label into an actionable surveillance state.

7. Clinical Translation of Biologic and Physiologic Profiling

A conceptual framework for biologic and physiologic signatures has limited value unless it can be translated into bedside decision support without becoming either opaque or clinically disruptive. Translation begins with a simple recognition: no clinician needs another isolated number detached from physiology. What is needed is a representation of leak risk that is interpretable, temporally updated, and clearly connected to possible clinical responses. The ideal output is not merely a probability but a physiologic synopsis of why the patient's trajectory is deviating and which mechanisms appear to dominate that deviation.

For this reason, the translational unit should probably not be a single scalar score displayed once per day. A more useful format would pair a calibrated probability with a dominant-signature profile. For example, the system might indicate that current risk is being driven primarily by delayed perfusion recovery and persistent inflammatory activation in the setting of low synthetic reserve. Such a profile does not claim mechanistic certainty. Rather, it provides a structured interpretation of the latent-state estimate [22]. This supports clinician trust because it maps the model output onto recognizable physiologic reasoning. In high-stakes surgical settings, the interpretive layer may matter as much as the numerical layer.

Translation also requires attention to measurement feasibility. Many biologically attractive variables are not routinely available. Direct tissue oxygen measurements, molecular markers of matrix turnover, or local microbiome states would almost certainly enrich leak-risk estimation, but they are not yet practical for universal deployment. A translational strategy should therefore distinguish core variables from extension variables. Core variables are routinely measured across institutions: inflammatory counts, lactate,

hemoglobin, basic metabolic indices, albumin or related reserve measures when available, hemodynamic summaries, vasopressor exposure, fluid balance, and procedure-specific context. Extension variables include advanced perfusion imaging, fluorescence-derived microvascular curves, wearable physiologic data, cytokine panels, and omic profiles. The core set must support useful inference on its own. The extension set should refine latent phenotyping when available, not determine whether the framework can function at all.

Calibration is a central translational requirement. A risk estimate is clinically meaningful only if its scale corresponds to observed frequencies under comparable conditions. Poorly calibrated models can rank patients reasonably well while misleading clinicians about absolute risk. This matters because the response to a 10% risk estimate is not the same as the response to a 40% estimate, even if the ordinal ranking is unchanged. Calibration becomes harder in leak surveillance because base rates vary across operation type, institutional practice, diversion strategies, and case mix. Therefore, a deployable system should include local recalibration layers or hierarchical priors that adjust absolute risk while preserving latent-state interpretation. Without such adaptation, a technically elegant model may fail precisely where clinicians most need reliability.

Clinical translation also depends on the intervention model attached to the risk signal. A model without a defined response pathway tends to generate either alarm fatigue or neglect. The response should be proportionate and state specific. A low-to-moderate but rising risk signal might prompt additional clinical review and closer physiologic monitoring. A high risk signal dominated by perfusion-related instability could trigger evaluation of volume status, vasopressor burden, or perfusion imaging where available. A high risk signal with inflammatory and barrier features may justify earlier imaging or source-control consideration. This state-specific logic is one reason signature profiling is preferable to generic scoring. It aligns surveillance with potential action.

Importantly, the value of a signature-based framework is not confined to diagnosis of established leaks. It can inform perioperative planning. Preoperative low-reserve signatures, such as depleted protein state, inflammatory activation, frailty-associated physiologic dampening, or poor vascular reserve, may indicate that the postoperative trajectory is likely to be narrow and easily destabilized [23]. Intraoperatively, such information may influence the threshold for diversion, the aggressiveness of hemodynamic optimization, or the willingness to accept marginal perfusion. Postoperatively, it may justify earlier surveillance in patients whose expected recovery capacity is limited even before obvious instability develops. Translation, therefore, spans the entire perioperative continuum.

A further issue is domain shift. The physiologic meaning of variables can differ across centers because monitoring intensity, resuscitation practices, surgical technique, critical care thresholds, and documentation conventions all vary. A variable such as ICU admission may carry strong signal in one environment and much weaker signal in another. Even laboratory timing changes variable interpretation. This means that deployable systems must be designed for transportability, not just internal performance. One approach is to emphasize mechanism-linked variables whose biological meaning is more stable than local workflow markers. Another is to separate universal latent dimensions from institution-specific observation models, allowing the same hidden-state framework to express itself differently depending on local data collection patterns.

Fairness and subgroup robustness are equally important. A biologic signature framework should not assume that identical numeric patterns mean identical risk across all patient populations without verification. Baseline inflammatory patterns, nutritional distributions, body composition, hemodynamic reserve, and access to perioperative optimization can differ systematically across demographic and clinical groups. If a model is trained on a narrow population, the latent space itself may be distorted, causing some groups to appear anomalous simply because the model has not learned their normal recovery geometry. Translation therefore requires explicit subgroup calibration and performance analysis. This is not an optional ethical add-on. It is a technical condition for valid inference.

The interpretability of signature-based profiling can be enhanced through graphical summaries. A trajectory display showing expected versus observed recovery along latent inflammatory, perfusion, and reserve dimensions may be more useful than a table of laboratory values. Surgeons and intensivists are

accustomed to integrating trajectories visually. A model that summarizes where the patient sits relative to typical healing paths may therefore integrate more naturally into practice than one that provides only a fixed threshold alert. Such visualization should remain parsimonious. The goal is not to overwhelm the clinician with model internals, but to externalize the core structure of the risk estimate.

Implementation also depends on timing within workflow. A biologic and physiologic profiling system would ideally operate at several moments: before surgery, immediately after surgery, during the first postoperative day, and at each major physiologic update thereafter. The preoperative estimate reflects reserve and baseline vulnerability. The early postoperative estimate incorporates the host response to operative stress. Later updates focus increasingly on whether recovery is organizing or failing [24]. This staged design acknowledges that risk is not static. It also prevents overreliance on either preoperative or postoperative data alone. Some patients are biologically fragile but recover well. Others appear robust but mount a maladaptive response to intraoperative insult. Translation requires seeing both possibilities.

The educational consequence of clinical deployment should not be underestimated. Signature-based systems can refine clinician intuition by making latent patterns explicit. Over time, this may improve bedside assessment even independent of the model. For example, teams may become more attentive to the joint significance of delayed lactate recovery, inflammatory persistence, and low reserve rather than treating each in isolation. In that sense, the model acts partly as a cognitive scaffold. Good translation, therefore, enhances clinical reasoning rather than attempting to replace it.

There is also a regulatory and methodological implication. Evaluation of such systems should not be limited to area under the curve or similar aggregate measures. A clinically mature assessment should examine lead time to actionable warning, stability of risk under repeated updates, calibration across subgroups, interpretability judged by clinician users, and the effect of model-guided surveillance on downstream practice. The most informative future studies will likely combine observational performance with pragmatic implementation testing. Only then can one determine whether biologic and physiologic profiling improves recognition, reduces delayed diagnosis, or alters operative and postoperative management in ways that matter.

Ultimately, the translational promise of this framework lies in reframing the anastomosis as a monitored healing system rather than a static technical accomplishment. The patient should not be judged solely by whether a leak has already declared itself. The more meaningful question is whether the signatures of repair remain coherent. If they do not, the clinical team gains an opportunity to intervene earlier, interpret postoperative uncertainty more rigorously, and move from reactive diagnosis toward physiologically informed surveillance.

8. Discussion

The central argument of this paper is that leak risk should be understood as the expression of a destabilized healing system rather than the consequence of a single abnormal variable or a purely technical failure. This reframing has several implications. It suggests that the most informative predictors are not isolated measurements but multivariate signatures linking tissue biology to systemic physiology. It also implies that risk estimation must be dynamic, because the transition from adaptive stress to structural failure unfolds in time. Finally, it indicates that useful models will need to combine mechanistic plausibility with statistical flexibility, especially when the available clinical data are incomplete and noisy.

One of the advantages of the signature framework is that it resolves a persistent tension in postoperative care. Clinicians know that many laboratory and hemodynamic abnormalities are nonspecific. Yet they also know that patients who later develop leaks often look subtly abnormal before the diagnosis becomes evident [25]. The gap between these observations arises because the bedside usually presents variables one at a time, while the biologic process acts through coupled patterns. A mild leukocytosis, a modest lactate elevation, and delayed physiologic normalization may each be unconvincing alone. When they co-occur in a patient with limited reserve and a structurally demanding anastomosis, the pattern becomes more meaningful. Signature analysis provides a formal language for this intuition.

A further insight is that leak risk is heterogeneous not only in magnitude but also in mechanism. Some patients likely fail along an ischemic path, where perfusion deficit and local necrosis dominate. Others fail along an inflammatory or septic path, in which microbial-barrier disruption and uncontrolled host response drive tissue breakdown. Others may fail because baseline reserve is insufficient for the catabolic and mechanical demands of healing. This heterogeneity helps explain why prediction has proven difficult when models rely on broad patient factors alone. If several mechanistic routes converge on the same endpoint, then a single global threshold may obscure important internal structure. A latent-state approach is attractive because it allows those routes to be represented as neighboring but distinct regions within the same risk manifold.

The biologic plausibility of this perspective is strong. Anastomotic healing requires simultaneous success in perfusion, inflammatory regulation, matrix reconstruction, and barrier restitution. The system is resilient to modest perturbation in one domain if the others compensate. It becomes fragile when compensatory coupling weakens. This is precisely the pattern that dynamic systems theory would predict: multi-component systems often fail not when one variable deviates slightly, but when the relationships that stabilize the variables become uncoupled. In clinical terms, the dangerous state is one in which inflammation persists while perfusion lags, reserve is depleted, and organ-level compensation remains burdened. The signature lies in the persistence of discordance.

The mathematical formulations proposed here are intentionally abstract, but they serve several purposes. They show how covariance can reveal dominant latent dimensions. They demonstrate how temporal evolution can be represented through state transitions and hazard accumulation. They clarify why interaction and nonlinearity are not optional embellishments but reflections of the underlying physiology. They also illustrate how patient monitoring can be reframed from threshold detection to state inference. Even simple versions of these models can improve conceptual rigor by forcing explicit assumptions about which variables are expected to cohere during normal healing and which deviations signal instability.

There are, however, important practical constraints. Routine clinical data are imperfect proxies for local tissue biology. Lactate is not a direct measure of anastomotic oxygenation. Leukocytosis is not specific to barrier failure [26]. Albumin is a noisy summary of reserve. Hemodynamic markers are shaped by resuscitation strategy as much as by intrinsic physiology. This means that no latent-state estimate should be confused with direct mechanistic observation. The proposed framework is best viewed as an inference architecture that maps imperfect measurements onto a probability distribution over biologically plausible states. Its strength lies in integration and updating, not in claiming molecular certainty from bedside data.

Another limitation is that signatures themselves can be context dependent. The same physiologic pattern may have different implications after a low colorectal anastomosis than after a proximal enteric reconstruction. Emergent surgery, diversion status, intraoperative contamination, steroid exposure, and baseline frailty all shift the meaning of subsequent measurements. This argues for hierarchical models in which procedure-specific and context-specific effects modulate the interpretation of shared latent dimensions. It also argues against transporting models without careful recalibration. A biologic concept may be universal while its observed numerical expression remains local.

The translational challenge is to preserve complexity without sacrificing usability. Surgeons and intensivists do not need exhaustive model internals. They need outputs that explain the dominant source of risk and indicate whether the trajectory is worsening, stabilizing, or improving. There is a danger that rich latent-state models become clinically inert if their output is opaque. Conversely, there is a danger that overly simplified bedside tools lose the very relational information that makes signatures useful. The best implementation will likely use complex internal estimation but simple external summaries, such as dominant risk axes, current trajectory direction, and lead-time calibrated probability.

An additional question concerns whether biologic and physiologic signatures can inform intervention, not just detection. The answer is likely yes, though with caution. If risk appears dominated by impaired perfusion and oxygen debt, clinicians may intensify efforts to optimize flow, reduce vasoconstrictive burden where feasible, and reassess local perfusion status. If risk is driven more by inflammatory

persistence and barrier stress, earlier imaging and source-control evaluation may be appropriate. If poor reserve dominates, nutritional and metabolic support become more salient, even if they cannot reverse imminent failure. The signature framework does not make treatment decisions automatically, but it can organize them according to mechanism-sensitive priorities.

This paper also implies a research agenda. Future observational work should focus less on accumulating isolated predictors and more on measuring coordinated trajectories. Datasets should preserve time resolution, intervention timing, and procedure context. Model development should incorporate explicit latent-state or manifold representations rather than relying entirely on static tabular abstraction. Validation should examine not only discrimination but lead time, calibration, subgroup robustness, and stability under repeated updating [27]. Mechanistic enrichment through microcirculatory imaging, perfusion fluorescence, molecular wound markers, and microbiome data should be pursued where feasible, but always anchored to a shared latent-state interpretation rather than treated as disconnected modalities.

There is also room for prospective physiologic studies aimed specifically at signature discovery. Instead of waiting for clinical leaks and retrospectively labeling them, investigators could serially profile the perioperative period to identify common trajectories of inflammatory resolution, perfusion recovery, and metabolic normalization. Patients who later develop leak could then be analyzed as departures from these normative trajectories. Such work would likely reveal that the earliest warning is often not an extreme value but a subtle failure of coordination among several moderate deviations.

In broader terms, the significance of the present framework extends beyond leak itself. Many postoperative complications arise from distributed failures of compensation and tissue repair. Anastomotic leak is a particularly consequential example because it sits at the intersection of microbiology, perfusion, matrix biology, and systemic stress. As such, it provides an ideal model for how modern perioperative medicine might move from event-based thinking to state-based surveillance. The biologic and physiologic signature approach is therefore relevant not only to colorectal or gastrointestinal surgery but also to the general problem of recognizing when a patient is no longer recovering in an organized way.

What remains essential is restraint in interpretation. No model can eliminate uncertainty from the postoperative period. Not every high-risk signature will culminate in leak, and not every leak will be heralded by a clean preclinical pattern. Biology is too heterogeneous and monitoring too incomplete for certainty. The purpose of signature-based reasoning is not to replace clinical judgment with deterministic classification. It is to make uncertainty more structured, earlier, and more biologically interpretable. That alone would represent meaningful progress.

9. Conclusion

Leak risk after gastrointestinal anastomosis is best understood as a multiscale instability of healing rather than as a purely technical mishap or the consequence of any single biomarker abnormality. The biologic substrate of this instability includes impaired microvascular oxygen delivery, dysregulated inflammatory control, defective matrix reconstruction, compromised epithelial barrier restitution, altered microbial containment, and inadequate metabolic reserve. The physiologic expression of the same process appears in patterns of delayed lactate normalization, persistent inflammatory activation, prolonged compensatory hemodynamic stress, organ-level burden, and failure of coordinated recovery.

A useful risk framework must therefore move beyond static variable lists and toward dynamic signature inference. In such a framework, laboratory values, vital sign trajectories, procedural context, and reserve markers are interpreted as correlated projections of latent states such as perfusion strain, inflammatory persistence, reparative insufficiency, and barrier vulnerability. Mathematical representations based on covariance geometry, state-space dynamics, and time-varying hazard provide one rigorous way to formalize these ideas. Their primary value lies not in elegance alone, but in their ability to align prediction with underlying physiology.

The translational implication is that perioperative surveillance should estimate where a patient lies within the manifold of organized recovery versus emerging instability, how rapidly that position is changing, and which physiologic axes are driving the change. This perspective supports earlier and

more interpretable warning, more coherent integration of multimodal data, and more mechanism-sensitive postoperative decision support. The anastomosis should thus be approached as a monitored healing system whose failure is preceded by detectable biologic and physiologic signatures. Making those signatures visible is the central technical challenge and the most plausible path toward more timely recognition of leak risk [28].

References

[1] A. Alakkari, R. Sood, S. M. Everett, B. J. Rembacken, J. D. Hayden, A. I. Sarela, and N. Mohammed, "First uk experience of endoscopic vacuum therapy for the management of oesophageal perforations and postoperative leaks," *Frontline gastroenterology*, vol. 10, pp. 200–203, 1 2019.

[2] J. Khoraki, G. S. Mazzini, A. S. Shah, P. A. R. D. Prado, L. G. Wolfe, and G. M. Campos, "Early small bowel obstruction after laparoscopic gastric bypass: A surgical emergency," *Surgery for obesity and related diseases : official journal of the American Society for Bariatric Surgery*, vol. 14, pp. 1118–1125, 5 2018.

[3] I. Pasternak, M. Dietrich, R. J. Woodman, U. Metzger, D. Wattchow, and U. Zingg, "Use of severity classification systems in the surgical decision-making process in emergency laparotomy for perforated diverticulitis.," *International journal of colorectal disease*, vol. 25, pp. 463–470, 11 2009.

[4] J. Orford, D. T. Cass, and M. J. Glasson, "Advances in the treatment of oesophageal atresia over three decades: the 1970s and the 1990s," *Pediatric surgery international*, vol. 20, pp. 402–407, 5 2004.

[5] C. J. Salgado, A. Nugent, J. Kuhn, M. G. Janette, and H. Bahna, "Primary sigmoid vaginoplasty in transwomen: Technique and outcomes," *BioMed research international*, vol. 2018, pp. 4907208–4907208, 5 2018.

[6] B. Sadanandan, "Data-driven risk stratification for anastomotic leak: A proof-of-concept study using electronic health records," *Journal of Data Science, Predictive Analytics, and Big Data Applications*, vol. 9, no. 6, pp. 1–27, 2024.

[7] B. Thakur, M. Devkota, N. Bhandari, S. Shrestha, and A. Kharel, "Minimally invasive esophagectomy/ gastroesophagectomy for cancer - long term results from a single institution," *Nepalese Journal of Cancer*, vol. 6, pp. 7–15, 10 2022.

[8] J. P. Burke, J. C. Coffey, E. Boyle, F. B. Keane, and D. A. McNamara, "Early outcomes for rectal cancer surgery in the republic of ireland following a national centralization program.," *Annals of surgical oncology*, vol. 20, pp. 3414–3421, 8 2013.

[9] S. K. Veetil, P. D. Haque, and D. Jain, "Clinical outcome analysis of immediate morbidity and mortality in patients presenting with obstructed colorectal cancers.," 4 2023.

[10] B. Flor-Lorente, J. F. Noguera-Aguilar, S. Delgado-Rivilla, J. M. García-González, M. Rodriguez-Martín, L. Salinas-Ortega, M. Ángel Casado, and M. Álvarez, "The economic impact of anastomotic leak after colorectal cancer surgery.," *Health economics review*, vol. 13, pp. 12–12, 2 2023.

[11] R. Mettke, A. Schmidt, S. Wolff, A. Koch, H. Ptok, H. Lippert, and I. Gastinger, "Spleen injuries during colorectal carcinoma surgery. effect on the early postoperative result," *Der Chirurg; Zeitschrift fur alle Gebiete der operativen Medizen*, vol. 83, pp. 809–814, 3 2012.

[12] P. Mowschenson, J. F. Critchlow, and M. A. Peppercorn, "Ileoanal pouch operation: long-term outcome with or without diverting ileostomy.," *Archives of surgery (Chicago, Ill. : 1960)*, vol. 135, pp. 463–466, 4 2000.

[13] M. El-Ahmar, F. Peters, M. Green, M. Dietrich, M. Ristig, L. Moikow, and J.-P. Ritz, "Robotic colorectal resection in combination with a multimodal enhanced recovery program - results of the first 100 cases.," *International journal of colorectal disease*, vol. 38, pp. 95–95, 4 2023.

[14] M. Hellan, J. R. Ouellette, J. A. Lagares-Garcia, S. M. Rauh, H. L. Kennedy, J. D. Nicholson, D. Nesbitt, C. S. Johnson, and A. Pigazzi, "Robotic rectal cancer resection: A retrospective multicenter analysis," *Annals of surgical oncology*, vol. 22, pp. 2151–2158, 12 2014.

[15] V. S. Budipramana and P. A. Saraswati, "Bishop-koop modification technique following proximal jejunal anastomosis: A case report.," *Annals of medicine and surgery (2012)*, vol. 56, pp. 121–124, 6 2020.

[16] V. Jayaraman, C. Hammerle, S. K. Lo, L. H. Jamil, and K. Gupta, "Clinical application and outcomes of over the scope clip device: Initial us experience in humans," *Diagnostic and therapeutic endoscopy*, vol. 2013, pp. 381873–381873, 7 2013.

[17] F. V. Mandarino, A. Barchi, L. Leone, L. Fanti, F. Azzolini, E. Viale, D. Esposito, N. Salmeri, F. Puccetti, L. Barbieri, A. Cossu, E. Treppiedi, U. Elmore, R. Rosati, and S. Danese, "Endoscopic vacuum therapy versus self-expandable metal stent for treatment of anastomotic leaks<30 mm following oncologic ivor-lewis esophagectomy: a matched case-control study.," *Surgical endoscopy*, vol. 37, pp. 7039–7050, 6 2023.

[18] L. Balanescu, A. Moga, R. Balanescu, T. Strimbu, and A. Cardoneanu, "Our experience with cyst excision and hepaticoenterostomy for choledocal cyst: A single center case review of 16 patients.," *Medicina (Kaunas, Lithuania)*, vol. 58, pp. 416–416, 3 2022.

[19] A. C. Chang, "Incisions and esophagectomy: is surgical approach all that matters?," *JAMA surgery*, vol. 148, pp. 739–739, 8 2013.

[20] A. Eroğlu, C. Daharlı, A. B. Ulaş, H. Keskin, and Y. Aydın, "Minimally invasive ivor-lewis esophagectomy for esophageal cancer.," *Turk gogus kalp damar cerrahisi dergisi*, vol. 30, pp. 421–430, 7 2022.

[21] C. Seifarth, J.-P. Ritz, U. Pohlen, A. J. Kroesen, B. Siegmund, B. Frericks, and H.-J. Buhr, "Therapy of complicated crohn's disease during pregnancy–an interdisciplinary challenge.," *International journal of colorectal disease*, vol. 29, pp. 645–651, 5 2014.

[22] U. S. Sibia, S. B. Badve, A. C. Istl, J. R. Klune, and A. I. Riker, "Impact of frailty upon surgical decision-making for left-sided colon cancer.," *Ochsner journal*, vol. 23, pp. 120–128, 4 2023.

[23] M. Rutegård and J. Rutegård, "Anastomotic leakage in rectal cancer surgery : The role of blood perfusion," *World journal of gastrointestinal surgery*, vol. 7, pp. 289–292, 11 2015.

[24] M. Senne, C. R. Werner, U. Schempf, K. Thiel, A. Königsrainer, and D. Wichmann, "Comparison of two endoscopic therapeutic interventions as primary treatment for anastomotic leakages after total gastrectomy.," *Cancers*, vol. 14, pp. 2982–2982, 6 2022.

[25] M. Fujita, K. Yamaguchi, K. Nagashima, K. Suzuki, T. Kasai, H. Hashimoto, Y. Onouchi, D. Sato, T. Fujisawa, and A. Hata, "Changes in colorectal cancer treatment during the covid-19 pandemic in japan: Interrupted time-series analysis using the national database of japan.," *Cancer epidemiology*, vol. 85, pp. 102391–102391, 5 2023.

[26] M.-P. Bernardi, A. G. Heriot, A. P. R. Sutherland, R. Woods, J. Keck, and M. Johnston, "Cr03 *ileocolic resections for crohn's disease: Impact of medical therapy," *ANZ Journal of Surgery*, vol. 79, 4 2009.

[27] M. Bhasin and P. K. Sachan, "Role of probiotics on surgical site infections in colorectal surgery," *International Surgery Journal*, vol. 7, pp. 1867–, 5 2020.

[28] J. Wu, L. R. Putnam, J. P. Silva, C. Houghton, N. Bildzukewicz, and J. C. Lipham, "Impact of robotic approach on post-anastomotic leaks after esophagectomy for esophageal cancer.," *The American surgeon*, vol. 88, pp. 2499–2507, 6 2022.